

METRONIDAZOLE & CLOTRIMAZOLE VAGINAL INSERT

DESCRIPTION:

Metronidazole & Clotrimazole Vaginal Insert is indicated for Nonspecific vaginitis, Bacterial vaginosis, Vulvo-vaginal candidiasis, Trichomonal vaginitis, vaginitis of mixed etiology, Dry vaginitis, recurrent vaginitis, infective leucorrhoea, Gram positive & Gram negative bacterial infections.

COMPOSITION:

Each vaginal inserts contains:

Metronidazole BP 500 mg Clotrimazole BP.....100 mg
ExcipientsQ.s.

DOSAGE:

Metronidazole & Clotrimazole Tablet should be inserted in the vaginal as deeply as possible. It should be used for 6 days, preferably on retiring to bed at night.

The treatment should not be carried out during menstruation.

SIDE EFFECT:

Vaginal burning, erythema, irritation & burning may rarely occur.

PRECAUTION:

Metronidazole & Clotrimazole Vaginal Insert should not be used if the patient has any of the following symptoms whereupon medical advice should be sought:

- Irregular vaginal bleeding.
- Abnormal vaginal bleeding or a blood-stained discharge.
- Vulval/vaginal ulcers, blisters or sores.
- Lower abdominal pain or dysuria.
- Fever or chills.
- Any adverse events such as redness, irritation or swelling associated with the treatment.
- Nausea or vomiting.
- Diarrhea.
- Foul smelling vaginal discharge.

PRESENTATION:

Metronidazole & Clotrimazole vaginal Insert are available in strip pack of 6 Tablets along with applicator.

